

## Heat-Related Illness Clinician's Guide June 2021

### Introduction

Agricultural workers are at significant risk for heat stress. Heat stress results when the body cannot get rid of excess heat and its core temperature rises.<sup>1</sup> Heat stress may lead to more severe heat illness including heat exhaustion, heat cramps, heat stroke, and even death if left untreated.<sup>2</sup> Agricultural work, which requires performing physically demanding work for long hours in hot and sometimes humid weather, places workers at high risk.

This guide provides information to clinicians on the prevention and treatment of heat-related illness. Since workers may not be familiar with all of the symptoms of heat stress, it is important that clinicians discuss heat illness symptoms and prevention with agricultural workers and others who are at risk.



© www.earldotter.com

### **Causes of heat-related illness**

High environmental temperatures, humidity, low or no-wind conditions, sun exposure, dehydration and physical exertion are all risk factors for heat illness.<sup>3</sup> Consuming alcohol or drinks high in caffeine—such as energy drinks—may increase the risk even further.<sup>4</sup> Personal factors such as age, weight, pregnancy, physical condition and use of certain medications may also put workers at greater risk (see *Importance of a thorough clinical history* on page 2). Serious heat illness is also more common among workers who are not accustomed to working in the heat. Approximately 50 to 70 percent of deaths attributed to outdoor heat exposure happen within the first few days of working in a warm or hot environment, due to lack of acclimatization.<sup>5</sup>

### **Recognizing and treating heat-related illness**

Since environmental and work conditions put agricultural workers at increased risk of heat illness, it is important that clinicians discuss this risk with patients and remain vigilant for symptoms that might indicate a patient is suffering from heat illness. Table 1 shows the different types of heat illness and symptoms.

Table 1. Types of heat illness and the second secon	heir symptoms
--	---------------

Heat stress	Heat exhaustion	Heat cramps	Heat stroke
<ul> <li>Feeling of discomfort</li> <li>Physiologic strain (indicated by increases in core temperature and heart rate in response to heat strain)</li> </ul>	<ul> <li>Rapid heartbeat</li> <li>Heavy sweating</li> <li>Extreme weakness or fatigue</li> <li>Dizziness</li> <li>Nausea, vomiting</li> <li>Irritability</li> <li>Fast, shallow breathing</li> <li>Slightly elevated body temperature</li> </ul>	• Muscle cramps, pain, or spasms in the abdomen, arms or legs	<ul> <li>High body temperature</li> <li>Confusion</li> <li>Loss of coordination (ataxia)</li> <li>Hot, dry skin or profuse sweating</li> <li>Throbbing headache</li> <li>Seizures, coma</li> </ul>

Sources: Jardine DS. Heat illness and heat stroke. *Pediatr Rev.* 2007 Jul;28(7):249-58. doi: 10.1542/pir.28-7-249. Erratum in: *Pediatr Rev.* 2007 Dec;28(12):469 and National Institute for Occupational Safety and Health. (2010). *NIOSH Fast Facts: Protecting Yourself from Heat Stress.* (DHHS (NIOSH) Publication No. 2010–114). https://www.cdc.gov/niosh/docs/2010-114/pdfs/2010-114.pdf.

### Chronic kidney disease of nontraditional origin (CKDnt):

Chronic heat stress may be associated with the development of CKDnt.<sup>6</sup> CKDnt is a form of kidney disease in patients who do not have the usual risk factors such as diabetes, hypertension, cardiovascular disease, or old age.<sup>7</sup> Some agricultural workers come to the U.S. from farms in Central America, where a CKDnt epidemic has killed tens of thousands of agricultural workers. The success of water-rest-shade interventions and the lack of CKDnt in subsistence farmers led researchers to conclude that occupational heat stress is the primary driver of the epidemic.<sup>8</sup> While rates of CKDnt in the U.S. are unknown, acute kidney injury (AKI) appears to be relatively common among agricultural workers.<sup>9</sup> Anecdotal information suggests that cases of illness consistent with CKDnt are increasingly being found in the U.S.<sup>10</sup>

CKDnt is typically asymptomatic except at advanced stages. Traditional clinical markers of kidney disease such as elevated serum creatinine, protein excretion and urinary albumin to creatinine ratio (ACR) may only develop when CKDnt is at an advanced stage, and there are currently no widely accepted reference standards for other biomarkers that might allow for early detection.<sup>11</sup> This underscores the importance of heat stress prevention among at-risk workers.

### Importance of a thorough clinical history:

Risk factors such as past episodes of heat illness and medical conditions or medications that affect the body's heat-regulation mechanism can make workers more susceptible to the effects of heat. Therefore, ascertaining a patient's clinical history is a key step toward identifying those at higher risk. Tables 2 and 3 show some physiological and pharmacological risk factors associated with a greater risk of heat illness.

# Table 2. Health conditions and individual factors that may increase workers' risk of heat-related illness

- Advanced age
- Altered cytokine production
- Cardiovascular disease
- Diabetes
- Hypohidrosis
- Kidney disease

- Low physical fitness
- Lung disease
- Malignant hyperthermia
- Overweight/obesity
- Pregnancy
- Previous episodes of heat illness

- Recent fever, diarrhea, cold
- Sickle cell trait
- Small body size
- Sunburn
- Sympathectomy
- Type II fiber predominance

Sources: Westwood CS, Fallowfield JL, Delves SK, Nunns M, Ogden HB, Layden JD. Individual risk factors associated with exertional heat illness: A systematic review. *Exp Physiol*. 2021 Jan;106(1):191-199. doi: 10.1113/EP088458. Epub 2020 May 8. Lundgren K, Kuklane K, Gao C, Holmér I. Effects of heat stress on working populations when facing climate change. Ind Health. 2013;51(1):3-15. doi: 10.2486/indhealth.2012-0089.

### Table 3. Medications/drugs that may increase workers' risk of heat-related illness

- Alcohol
- Alpha-adrenergic agonists
- Amphetamines
- Anticholinergics
- Antihistamines
- Benzodiazepines
- Vasoconstrictors including migraine and allergy medications
- Antipsychotic Drugs (ACDs)

- Beta blockers
- Calcium channel blockers
- Cocaine
- Diuretics
- Ephedra-containing supplements
- Laxatives
- Selective Serotonin Reuptake Inhibitors (SSRIs)

- Neuroleptics
- Phenothiazines
- Stimulants
- Thyroid receptor agonists
- Tricyclic antidepressants
- Serotonin Norepinephrine Reuptake Inhibitors (SNRIs)

This list of medications is not exhaustive. Certain medications in combination may also increase risk. See Resources for more information.

Source: Becker JA, Stewart LK. Heat-related illness. Am Fam Physician. 2011 Jun 1;83(11):1325-30. Serani D. Heat intolerance and psychiatric medications. Psychology Today. 27 July 2021.

### **Prevention:**

Drinking water frequently, taking shade and rest breaks, and allowing for a period of acclimatization to working in the heat are key steps to prevent heat illness. The OSHA field sanitation standard requires employers with more than 10 workers to provide drinking water.<sup>12</sup> It is recommended that workers drink 1 cup (8 oz.) of water or other fluids every 15-20 minutes when working in hot conditions. Agricultural workers should also take rest/shade breaks, which should increase in duration and frequency according to the temperature and the intensity of their work activity.<sup>13</sup> Other basic messages include to avoid alcohol and caffeinated drinks like soda or energy drinks and to opt for water instead; to wear clothing like loose-fitting, lightweight, light-colored cotton clothes, a wide-brimmed hat, and a bandana; and to monitor oneself and coworkers closely during heat events.



Agricultural workers may be paid by piece rate rather than by the hour, disincentivizing workers from taking time to get water and take rest/shade breaks. Clinicians should emphasize the importance of close monitoring of signs and symptoms in addition to strongly advocating for water, rest, and shade. Clinicians may also contact employers to encourage worker and supervisor training to prevent heat-related deaths.

### **Treatment and medical attention:**

If the worker has symptoms of heat stress or heat exhaustion, he/she should be removed from the heat immediately, taken to a shaded, cool place and given fluids to drink. If the symptoms continue after 20 or 30 minutes, emergency medical attention may be needed. Any decline in mentation, regardless of core temperature, must be considered heat stroke. Emergency medical attention is needed if a person is experiencing heat stroke symptoms such as ataxia, confusion, seizures or loss of consciousness.<sup>14</sup> While emergency medical services are summoned, the worker should be placed in a cool location, have excess clothing removed, and be wetted with cool water and fanned.<sup>15</sup> Once under medical care, cooling—through external and/or internal methods—must take priority and should continue until the patient's core temperature is reduced to 38 °C (100.4 °F). Once the cooling process is underway, laboratory tests and monitoring of renal function may be performed.<sup>16</sup>

### **Post-treatment follow-up:**

After counseling the worker on how to prevent the recurrence of heat illness, the clinician should provide the worker with a note for their employer indicating the reason the worker required medical treatment and any accommodations the employee may need upon their return to work. The U.S. Occupational Safety and Health Administration (OSHA) recommends that if a clinician determines that work conditions at a worksite are unsafe, they should discuss this with the employer while maintaining patient confidentiality. The clinician may also file a complaint with their state OSHA Office. If the conditions at a worksite are life-threatening, they should be reported immediately through OSHA's emergency number: 1-800-321-OSHA.

### Messaging to prevent heat-related illness:

While work conditions may not always provide an environment that is favorable to preventive practices, it is important that agricultural workers be aware of the need to drink water throughout the day and take regular shade and rest breaks without waiting to feel sick. The Centers for Disease Control and Prevention (CDC) has an <u>online resource</u> with answers to frequently asked questions about heat illness in English and Spanish.

# Social determinants of health and heat stress

Agricultural workers face specific social determinants of health (SDOH) that place them at greater risk of developing heat illness. SDOH commonly affecting agricultural workers include lack of access to clean water, lack of air conditioning at home to cool off and recover after work, and socioeconomic vulnerability and language barriers that may prevent them from demanding safer work conditions from their employer. They may also fear retaliation form their employer for denouncing unsafe work conditions.



### Other considerations around heat-related illness

There are additional factors to consider regarding heat-illness risks among agricultural workers. One of these is the amount of clothing and personal protective equipment (PPE) workers must wear while in the fields to protect themselves against COVID-19, dust inhalation, pesticide poisoning, and sunburn. PPE, although necessary, traps heat and can lead the body to overheat more quickly.

Current climate trends indicate that extreme heat events and average temperatures will continue to increase in the U.S. and with them the potential for heat illness. Another consequence of climate change is the increasing frequency of large wildfires. Even when burning hundreds of miles away from the agricultural worksite, wildfires can expose workers to the stress of wildfire smoke. In heavy smoke, many workers are encouraged to wear PPE to prevent smoke inhalation, increasing their risk of heat illness. Furthermore, heat stress may worsen the health effects of other environmental conditions such as air pollution.<sup>17</sup>

### Heat-related illness and the law

There is no national heat standard to protect agricultural workers from the risk of heat illness. As of this writing, only three states—California, Minnesota and Washington—have workplace heat standards. Due to the lack of a national occupational heat standard and imperfect compliance in states that do have regulations in place, many workers are not afforded the opportunity to take basic preventive measures against heat stress.<sup>18</sup>

A bill introduced in the 117<sup>th</sup> Congress in 2021—the "<u>Asunción Valdivia Heat Illness and</u> <u>Fatality Prevention Act</u>"—would require OSHA to create a national heat standard mandating paid rest breaks in cool or shaded environments for outdoor workers, access to drinking water and worker training on heat stress, among other provisions.<sup>19</sup> The bill was named after an agricultural worker who tragically died from heat stress. However, a bill is not needed for OSHA to take action at the agency level to issue a heat stress standard. More information on heat-related illness and pending legislation can be found on the Farmworker Justice website.

### **Clinician Resources**

- <u>Heat-Related Illnesses</u>, by R. Gauer and B.K. Meyers<sup>20</sup>
- Emergency Management of Heat-Related Illness, by W. Troyer and J. Kiel<sup>21</sup>
- <u>Heat Illness</u>, by D. Leiva and B. Church<sup>22</sup>
- Evaluation and Management of Heat Stroke, by K. Li<sup>23</sup>
- <u>Heatstroke</u>, by Y. Epstein and R. Yanovich<sup>24</sup>
- <u>Management of exertional heat stroke: a practical update for primary care</u> <u>physicians</u>, by E.Walter and K. Steel<sup>25</sup>
- <u>Management of Environmental Heat Injury in the ED</u>, by S. Watts, P. Jackson and G. Chiampas<sup>26</sup>
- <u>Management of Heatstroke and Heat Exhaustion</u>, by J.L. Glazer<sup>27</sup>

### **Patient Resources**

The following are some useful resources designed specifically for workers on how to prevent, recognize and address heat illness in the workplace.

#### **Centers for Disease Control and Prevention**

- <u>Heat-Related Illneses</u> (English)
- Enfermedades Relacionadas con el Calor (Spanish)

#### National Institute for Occupational Safety & Health

- <u>NIOSH Fast Facts</u> (English)
- <u>NIOSH Datos Breves</u> (Spanish)

#### **Occupational Safety and Health Administration**

- <u>Water. Rest. Shade. The Work Can't Get Done Without Them</u> (Low literacy fact sheet)
- Agua. Sombra. Descansos. Sin ellos no se puede trabajar (Spanish)

#### Pacific Northwest Agricultural Safety and Health Center

Heat Illness Prevention

#### Western Center for Agricultural Health and Safety

Heat Illness Prevention

#### Association of Farmworker Opportunity Programs

Heat Stress Prevention

### **Employer Resources**

Clinicians may share resources with local employers.

#### **Occupational Safety and Health Administration:** Using the Heat Index: <u>A Guide for Employers</u>

Agua. Sombra. Descansos. Sin ellos no se puede trabajar (Spanish)

#### California\* Occupational Safety and Health Administration:

<u>Heat Illness Prevention</u> Prevención de Enfermedades Causadas por el Calor

\*California requires employers to provide trainings on heat illness prevention. However, much of these materials – including trainings, posters, and a pocket guide, are applicable to employers around the country to train workers.

**For more information,** contact Iris Figueroa, Director of Economic and Environmental Justice, Farmworker Justice at <u>ifigueroa@farmworkerjustice.org</u> or Amy Liebman, Director of Environmental and Occupational Health, Migrant Clinicians Network, at <u>aliebman@migrantclinician.org</u>.

This joint FJ and MCN publication was supported by the Health Resources and Services Administration (HRSA) of the U.S. Department of Health and Human Services (HHS) as part of awards totaling \$1,949,598 with 0% financed with non-governmental sources. The contents are those of the authors and do not necessarily represent the official views of, nor an endorsement by, HRSA, HHS, or the U.S. Government. For more information, please visit HHS.gov.

### References

<sup>1</sup> Choudhary E, Vaidyanathan A. Heat Stress Illness Hospitalizations — Environmental Public Health Tracking Program, 20 States, 2001–2010. *Morbidity and Mortality Weekly Report* 2014; 63(SS13);1-10.

- https://www.cdc.gov/mmwr/preview/mmwrhtml/ss6313a1.htm.
- <sup>2</sup> National Institute for Occupational Safety and Health (NIOSH). (2018). Heat Stress Heat Related Illness.
- https://www.cdc.gov/niosh/topics/heatstress/heatrelillness.html.
- <sup>3</sup> OSHA. (2014). OSHA Fact Sheet: Protecting Workers from the Effects of Heat. (OSHA Publication No. DTSEM FS-3743 08/2014). https://www.osha.gov/sites/default/files/publications/heat\_stress.pdf.
- <sup>4</sup> NIOSH. (2017.) Heat Stress: Hydration. (DHHS (NIOSH) Publication No. 2017-126).
- https://www.cdc.gov/niosh/mining/UserFiles/works/pdfs/2017-126.pdf.

<sup>5</sup> OSHA. *Heat.* https://www.osha.gov/heat-exposure.

<sup>6</sup> Laws RL, Brooks DR, Amador JJ, Weiner DE, Kaufman JS, Ramirez-Rubio O, Riefkohl A, Scammell MK, López-Pilarte D, Sánchez JM, Parikh CR, McClean MD. Biomarkers of Kidney Injury Among Nicaraguan Sugarcane Workers. *Am J Kidney Dis* 2016;67:209–17.<u>doi:10.1053/j.ajkd.2015.08.022</u>; Nerbass FB, Pecoits-Filho R, Clark WF, Sontrop JM, McIntyre CW, Moist L. Occupational Heat Stress and Kidney Health: From Farms to Factories. Kidney Int Rep. 2017 Aug 31;2(6):998-1008. doi: 10.1016/j.ekir.2017.08.012.

<sup>7</sup> Crowe J, Wesseling C, Solano BR, Umaña MP, Ramírez AR, Kjellstrom T, Morales D, Nilsson M. Heat exposure in sugarcane harvesters in Costa Rica. *Am J Ind Med.* 2013 Oct;56(10):1157-64. doi: 10.1002/ajim.22204. Epub 2013 Jun 17.

<sup>8</sup> Wesseling, C., Glaser, J., Rodríguez-Guzmán, J., Weiss, I, Lucas, R, Peraza, S, da Silva, AS, Hansson, EJohnson, RJ, Hogstedt, C, Wegman, DH, & Jakobsson, K (2020). Chronic kidney disease of non-traditional origin in Mesoamerica: a disease primarily driven by occupational heat stress. *Revista panamericana de salud publica = Pan American journal of public health*, *44*, e15. https://doi.org/10.26633/RPSP.2020.15

<sup>9</sup> Moyce S, Mitchell D, Armitage T, Tancredi D, Joseph J, Schenker M. Heat strain, volume depletion and kidney function in California agricultural workers. *Occup Environ Med.* 2017 Jun;74(6):402-409. doi: 10.1136/oemed-2016-103848. Epub 2017 Jan 16. Erratum in: Occup Environ Med. 2018 Feb;75(2):162; Mix J, Elon L, Vi Thien Mac V, Flocks J, Economos E, Tovar-Aguilar AJ, Stover Hertzberg V, McCauley, LA. Hydration Status, Kidney Function, and Kidney Injury in Florida Agricultural Workers. J Occup Environ Med. 2018 May;60(5):e253-e260. doi: 10.1097/JOM.00000000001261.

<sup>10</sup> Fernandez E. On the trail of a deadly disease that cuts down farmworkers in their prime. *Bloomberg Businessweek* 2018; Sep 25. https://www.bloomberg.com/news/articles/2018-09-25/ckdu-disease-that-kills-sugar-workers-is-spreading-in-the-u-s.
 <sup>11</sup> Gunasekara TDKSC, De Silva PMCS, Herath C, Siribaddana S, Siribaddana N, Jayasumana C, Jayasinghe S, Cardenas-Gonzalez M, Jayasundara N. The Utility of Novel Renal Biomarkers in Assessment of Chronic Kidney Disease of Unknown Etiology (CKDu): A Review. *Int J Environ Res Public Health*. 2020 Dec 18;17(24):9522. doi: 10.3390/ijerph17249522.

<sup>12</sup> OSHA Field Sanitation Standard. 29 CFR §1928.110. https://www.ecfr.gov/cgi-bin/text-

idx?SID=0ccc5d2974134cf6b65166500a7c417f&mc=true&node=se29.9.1928 1110&rgn=div8

<sup>13</sup> NIOSH. (2016). *Criteria for a Recommended Standard: Occupational Exposure to Heat and Hot Environments*. (DHHS (NIOSH) Publication 2016-106.) https://www.cdc.gov/niosh/docs/2016-106/pdfs/2016-

106.pdf?id=10.26616/NIOSHPUB2016106.

<sup>14</sup> Glazer JL. Management of heatstroke and heat exhaustion. *Am Fam Physician*. 2005 Jun 1;71(11):2133-40. PMID: 15952443. https://www.aafp.org/afp/2005/0601/afp20050601p2133.pdf.

<sup>15</sup> NIOSH. (2013). *Preventing Heat-related Illness or Death of Outdoor Workers*. (DHHS (NIOSH) Publication No. 2013–143). https://www.cdc.gov/niosh/docs/wp-solutions/2013-143/pdfs/2013-143.pdf?id=10.26616/NIOSHPUB2013143.

<sup>16</sup> Glazer JL. Management of heatstroke and heat exhaustion. *Am Fam Physician*. 2005 Jun 1;71(11):2133-40. PMID: 15952443. https://www.aafp.org/afp/2005/0601/afp20050601p2133.pdf. <sup>17</sup> De Sario M, Katsouyanni K, Michelozzi P. Climate change, extreme weather events, air pollution and respiratory health in Europe. *Eur Respir J.* 2013 Sep;42(3):826-43. doi: 10.1183/09031936.00074712. Epub 2013 Jan 11.

<sup>18</sup> State of California, Department of Industrial Relations. *California Code of Regulations, Title 8, section 3395. Heat Illness Prevention in Outdoor Places of Employment.* https://www.dir.ca.gov/title8/3395.html.

Minnesota Legislature. *Minnesota Administrative Rules.* 5205.0110 Indoor Ventilation and Temperature in Places of Employment. https://www.revisor.mn.gov/rules/5205.0110/.

Washington State Legislature. Outdoor Heat Exposure. WAC 296-307-097 et seq.

https://app.leg.wa.gov/WAC/default.aspx?cite=296-307.

<sup>19</sup> 117<sup>th</sup> Congress. S.1068 - Asuncion Valdivia Heat Illness and Fatality Prevention Act of 2021.

https://www.congress.gov/bill/117th-congress/senate-bill/1068/text?r=24&s=1

<sup>20</sup> Gauer R, Meyers BK. Heat-Related Illnesses. Am Fam Physician. 2019 Apr 15;99(8):482-489.

<sup>21</sup> Troyer W, Kiel J. Emergency Management of Heat-Related Illness. *EMResident*. 2018 Oct 16.

https://www.emra.org/emresident/article/heat-related-illness/

<sup>22</sup> Leiva DF, Church B. Heat Illness. [Updated 2021 Apr 15]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2021 Jan-. https://www.ncbi.nlm.nih.gov/books/NBK553117/

<sup>23</sup> Li, K. Evaluation and Management of Heat Stroke. ALiEM, 2015 Oct 28. https://www.aliem.com/heat-stroke

<sup>24</sup> Epstein Y, Yanovich R. Heatstroke. N Engl J Med 2019; 380:2449-2459. doi: 10.1056/NEJMra1810762

<sup>25</sup> Walter E, Steel K. Management of exertional heat stroke: a practical update for primary care physicians. Br J Gen Pract. 2018;68(668):153-154. doi:10.3399/bjgp18X695273

<sup>26</sup> Watts S, Jackson P. (2020, July 6). Management of Environmental Heat Injury in the ED. *NUEM Blog*, 2020 Jul 6. [Expert Commentary by Chiampas G.] http://www.nuemblog.com/blog/environmental-heat-injury

<sup>27</sup> Glazer JL. Management of heatstroke and heat exhaustion. *Am Fam Physician*. 2005 Jun 1;71(11):2133-40. PMID: 15952443. https://www.aafp.org/afp/2005/0601/afp20050601p2133.pdf.